

Implications of precision measurements on texture specific fermion mass matrices

Manmohan Gupta¹, Gulsheen Ahuja¹, Rohit Verma²

¹*Department of Physics, Centre of Advanced Study, P.U., Chandigarh, India.*

²*Rayat Institute of Engineering and Information Technology, Ropar, India.*

Email: mmgupta@pu.ac.in

November 4, 2009

Abstract

Implications of texture specific mass matrices have been investigated for both quarks and neutrinos. Interestingly, for the case of quarks Fritzsch-like texture 4 zero mass matrices have been found to be compatible with the present precisely known $\sin 2\beta$ as well as other precise CKM matrix elements. In the case of leptonic mass matrices, for both Majorana and Dirac neutrinos we find that for texture 4, 5, 6 zero mass matrices the inverted hierarchy and degenerate scenarios of neutrino masses are ruled out by the present data.

Understanding fermion masses and mixings constitutes one of the most important problems of flavor physics. The problem becomes mysterious when one finds that in the case of quarks the mixing angles as well as the quark masses show distinct hierarchy, this being in sharp contrast to the case of neutrinos wherein neither the mixing angles nor the masses follow a well defined hierarchy. In particular, at present there is no consensus about neutrino masses which may show normal/inverted hierarchy or may even be degenerate. In this context, Fritzsch-like texture specific mass matrices [1, 2] have given valuable insight in understanding several features of fermion masses and mixings. Usually the elements of the quark mass matrices are assumed to be hierarchical to explain the strongly hierarchical mixing angles, in contrast to the case of neutrinos where neutrino mixing parameters do not enforce any hierarchy on the elements of the mass matrices.

Several parameters in the case of quarks and leptons are now known with good deal of precision. In the case of quarks, the parameter $\sin 2\beta$ and the Cabibbo-Kobayashi-Maskawa (CKM) [3] matrix elements V_{us}, V_{cb}, V_{ud} are known precisely [4], however, V_{ub} is not that precisely known. In this context, our recent study [5] indicates that $\sin 2\beta$ plays a crucial role in fixing V_{ub} . Similar to the case of quarks, in the case of neutrinos angles s_{12}, s_{23} as well as the mass square differences $\Delta m_{12}^2, \Delta m_{23}^2$ are now rather well

known [4, 6], whereas only an upper limit is known regarding the third mixing angle s_{13} . Further, at present not much is known about the hierarchy of neutrino masses as well as about their absolute values.

The purpose here, on the one hand, is to check the compatibility of texture 4 zero mass matrices with the latest precisely known CKM parameters, in particular with $\sin 2\beta$. On the other hand, one would like to examine the implications of the present knowledge of neutrino oscillation parameters on the hierarchy of neutrino masses in the context of texture specific mass matrices.

Before we examine the compatibility of texture 4 zero mass matrices with the latest data, we would first like to briefly discuss the implications of $\sin 2\beta$, along with the unitarity of the CKM matrix, on the CKM element V_{ub} . Considering the usual ‘ db ’ unitarity triangle and definitions of the angles of the unitarity triangle, one arrives at a relation [5] involving CP violating phase δ , the mixing angles s_{12} , s_{23} , s_{13} as well as the angle β of the unitarity triangle, e.g.,

$$\tan \frac{\delta}{2} = \frac{A - \sqrt{A^2 - (B^2 - A^2 C^2) \tan^2 \beta}}{(B + AC) \tan \beta}, \quad (1)$$

where $A = c_{12}s_{12}s_{13}$, $B = c_{23}s_{23}(s_{12}^2 - c_{12}^2 s_{13}^2)$ and $C = c_{23}^2 - s_{23}^2$. Using $s_{12}^2 \gg c_{12}^2 s_{13}^2$ and $s_{23}^2 \ll c_{23}^2$, the above relation can be re-expressed as

$$\delta = -\beta + \sin^{-1} \left(\frac{s_{12}s_{23}}{c_{12}s_{13}} \sin \beta \right). \quad (2)$$

Further, using the closure property of the angles of the triangle, the mixing angle s_{13} can be written as

$$s_{13} = \frac{s_{12}s_{23} \sin \beta}{c_{12} \sin \alpha}. \quad (3)$$

Using these unitarity based relations we obtain $V_{ub} > 0.00355$, which on using the recently measured angle α of the unitarity triangle, translates to $V_{ub} = 0.0035 \pm 0.0002$. For our subsequent analysis, we will be using this value of V_{ub} .

After fixing V_{ub} , we would now like to examine, in detail, the compatibility of texture 4 zero mass matrices with the present value of $\sin 2\beta$ as well as with the other precisely known CKM elements. To this end, it may be noted that realizing the importance of the parameter $\sin 2\beta$ for the texture specific mass matrices, several authors [7]-[9] have explored its implications on these. In particular, using assumption of strong hierarchy of the elements of the mass matrix, they arrive at leading order relationships between the various elements of the mixing matrices and the quark masses. Further, using these relations and following PDG definition, in the strong hierarchy case, the angle β of the unitarity triangle comes out to be

$$\beta \equiv \arg \left[-\frac{V_{cd}V_{cb}^*}{V_{td}V_{tb}^*} \right] = \arg \left[1 - \sqrt{\frac{m_u m_s}{m_c m_d}} e^{-i\phi} \right]. \quad (4)$$

Unfortunately, the value of $\sin 2\beta$ predicted by the above formula is in quite disagreement with its present precisely known value. In particular, with the present values of

input quark masses, the value of $\sin 2\beta$ comes out to be < 0.45 , which is in sharp conflict with its present value given by PDG 2008 [4], e.g., 0.681 ± 0.025 . This is to be contrasted with some recent analyses [10, 11] which do not indicate this anomaly. This immediately brings forth the issues of modification of the above formula as well as of using strongly hierarchical mass matrices in the context of quarks. In this context, it may be noted that in principle it is an easy task to find an exact expression for β in the case of texture 4 zero mass matrices, however such an expression does not yield any useful information as it involves 4 V_{CKM} elements, each of which has complicated dependence on quark masses, phases and free parameters of quark mass matrices. Therefore, the first step in this direction is to develop a formula for $\sin 2\beta$ which allows one to go beyond the leading order, incorporate the correct phase structure of the mass matrices as well as allows one to understand the implications of the hierarchy on the mass matrices.

To facilitate the understanding of the relationship of the present work with the earlier attempts, we first define the modified Fritzsch-like matrices, e.g.,

$$M_i = \begin{pmatrix} 0 & A_i & 0 \\ A_i^* & D_i & B_i \\ 0 & B_i^* & C_i \end{pmatrix}, \quad i = U, D, \quad (5)$$

M_U and M_D , respectively corresponding to the mass matrix in the up sector and the down sector. It may be noted that each of the above matrix is texture 2 zero type with $A_i = |A_i|e^{i\alpha_i}$ and $B_i = |B_i|e^{i\beta_i}$.

Without getting into the details, using only the hierarchy of quark masses, $m_t \gg m_u$ and $m_b \gg m_d$, one can express β in terms of the quark masses as well as the phases of the quark mass matrix, e.g.,

$$\beta \equiv \arg \left[-\frac{V_{cd}V_{cb}^*}{V_{td}V_{tb}^*} \right] = \arg \left[\left(1 - \sqrt{\frac{m_u m_s}{m_c m_d}} e^{-i(\phi_1 + \phi_2)} \right) \left(\frac{1 - r_2 e^{i\phi_2}}{1 - r_1 e^{i\phi_2}} \right) \right], \quad (6)$$

wherein

$$r_1 = \frac{\zeta_{1U} \zeta_{3U} \zeta_{1D} \zeta_{2D}}{\zeta_{2U} \zeta_{3D}} \quad \text{and} \quad r_2 = \frac{\zeta_{1U} \zeta_{3U} \zeta_{2D}}{\zeta_{2U} \zeta_{1D} \zeta_{3D}}, \quad (7)$$

with ζ_{1i} , ζ_{2i} , ζ_{3i} (i denoting U and D) given by

$$\zeta_{1i} = \sqrt{1 + \frac{m_2}{C_i}}, \quad \zeta_{2i} = \sqrt{1 - \frac{C_i}{m_3}}, \quad \zeta_{3i} = \sqrt{\frac{C_i}{m_3}}. \quad (8)$$

The two phases ϕ_1 and ϕ_2 are defined as

$$\phi_1 = \alpha_U - \alpha_D \quad \text{and} \quad \phi_2 = \beta_U - \beta_D. \quad (9)$$

It may be noted that the relationship of β derived by us, given in equation (6), is almost an exact formula emanating from texture 4 zero mass matrices, incorporating both the phases ϕ_1 and ϕ_2 as well as no restriction on the hierarchy of the elements of the mass matrices.

Before discussing the results of our analysis, we would first like to briefly mention the inputs used for carrying out the calculations. We have adopted the following ranges of quark masses [12] at the energy scale of M_z , e.g.,

$$\begin{aligned} m_u &= 0.8 - 1.8 \text{ MeV}, & m_d &= 1.7 - 4.2 \text{ MeV}, & m_s &= 40.0 - 71.0 \text{ MeV}, \\ m_c &= 0.63 \text{ GeV}, & m_b &= 2.9 \text{ GeV}, & m_t &= 172.5 \text{ GeV}. \end{aligned} \quad (10)$$

Further, we have given full variation to the phases ϕ_1 and ϕ_2 , the parameters D_U and D_D have respectively been varied from 0 to $(m_t - m_c)$ and 0 to $(m_b - m_s)$. Furthermore, we have used the constraints imposed by the following well known CKM parameters [4],

$$\begin{aligned} |V_{us}| &= 0.2255 \pm 0.0019, & |V_{cb}| &= (41.2 \pm 1.1)10^{-3}, & |V_{ub}| &= 0.0035 \pm 0.0002, \\ \sin 2\beta &= 0.681 \pm 0.025. \end{aligned} \quad (11)$$

As mentioned earlier, we have considered the value of $|V_{ub}|$ obtained recently [5] using only the unitarity of the elements of the CKM matrix and current $\sin 2\beta$ value.

As a first step of our analysis, using these inputs we have obtained the CKM matrix at 1σ C.L. as follows

$$V_{\text{CKM}} = \begin{pmatrix} 0.9738 - 0.9747 & 0.2236 - 0.2274 & 0.0033 - 0.0037 \\ 0.2234 - 0.2273 & 0.9729 - 0.9739 & 0.0401 - 0.0423 \\ 0.0067 - 0.0103 & 0.0390 - 0.0418 & 0.9991 - 0.9992 \end{pmatrix}. \quad (12)$$

A general look at the matrix reveals that the ranges of CKM elements obtained here are quite compatible with those obtained by recent global analyses [4, 13, 14, 15].

After having checked the compatibility of texture 4 zero mass matrices with the latest CKM matrix elements, we would now like to briefly discuss the implications of compatibility of $\sin 2\beta$ on these matrices. Our calculations indicate that $\sin 2\beta$ has important implications for the hierarchy as well as the phase structure of the mass matrices. It is interesting to note that if we consider the particular case of strong hierarchy of the elements of the mass matrices, then even if we use the exact formula, derived here, given in equation (6), we are not able to reproduce $\sin 2\beta$ within its experimental limits. This clearly indicates that hierarchy of mass matrices play an important role in fitting the data. Further, despite giving full variation to other parameter and using the exact formula, in case we consider $\phi_2 = 0^\circ$, again we are not able to fit $\sin 2\beta$. These conclusions can be well understood by comparing the relationship derived by us, given in equation (6), with the earlier used leading order expression, given in equation (4). Now, the formula has an additional term $\left(\frac{1-r_2 e^{i\phi_2}}{1-r_1 e^{i\phi_2}}\right)$ which leads to significant contribution to $\sin 2\beta$. This happens only in case when $\phi_2 \neq 0^\circ$ and D_U/C_U or $D_D/C_D > 0.07$, implying somewhat weak hierarchy of the elements of the mass matrices.

As a next step of our analysis we have discussed the implications of the present knowledge of neutrino oscillation parameters on the texture structure of neutrino mass matrices, examined by us [16, 17, 18]. In particular, we have studied the issues pertaining to hierarchy of neutrino masses, e.g., the normal hierarchy case is defined as $m_{\nu_1} < m_{\nu_2} \ll m_{\nu_3}$, the

inverted hierarchy case is given by $m_{\nu_3} \ll m_{\nu_1} < m_{\nu_2}$ and the corresponding degenerate cases are respectively defined as $m_{\nu_1} \lesssim m_{\nu_2} \sim m_{\nu_3}$ and $m_{\nu_3} \sim m_{\nu_1} \lesssim m_{\nu_2}$. For both Majorana as well as Dirac neutrinos, we have imposed Fritzsch-like texture structure on Dirac neutrino mass matrix, with charged leptons having either Fritzsch-like texture structure or being in the flavor basis.

Considering the modified Fritzsch-like matrices, e.g.,

$$M_l = \begin{pmatrix} 0 & A_l & 0 \\ A_l^* & D_l & B_l \\ 0 & B_l^* & C_l \end{pmatrix}, \quad M_{\nu D} = \begin{pmatrix} 0 & A_\nu & 0 \\ A_\nu^* & D_\nu & B_\nu \\ 0 & B_\nu^* & C_\nu \end{pmatrix}, \quad (13)$$

M_l and $M_{\nu D}$ respectively corresponding to Dirac-like charged lepton and neutrino mass matrices, we have investigated 42 distinct possibilities of texture 4 zero, 5 zero and 6 zero mass matrices for normal/inverted hierarchy as well as degenerate scenarios of neutrino masses. The texture 6 zero matrices correspond to the above mentioned matrices when both D_l and D_ν are zero whereas texture 5 zero matrices correspond to either $D_l = 0$ and $D_\nu \neq 0$ or $D_\nu = 0$ and $D_l \neq 0$, the above matrices are texture 4 zero when both D_l and D_ν are non-zero.

The latest situation regarding masses and mixing angles at 3σ C.L. is summarized as follows [6],

$$\Delta m_{12}^2 = (7.14 - 8.19) \times 10^{-5} \text{ eV}^2, \quad \Delta m_{23}^2 = (2.06 - 2.81) \times 10^{-3} \text{ eV}^2, \quad (14)$$

$$\sin^2 \theta_{12} = 0.263 - 0.375, \quad \sin^2 \theta_{23} = 0.331 - 0.644, \quad \sin^2 \theta_{13} \leq 0.046. \quad (15)$$

To begin with, for Fritzsch-like texture 4, 5 and 6 zero lepton mass matrices detailed predictions for cases pertaining to inverted hierarchy as well as degenerate scenarios of neutrino masses have been carried out. For Majorana as well as Dirac neutrinos, our rigorous analysis, wherein all the input parameters have been given 3σ variations, reveals that all these cases seem to be ruled out. For the present purpose, we illustrate the case of texture 4 zero mass matrices. In figures 1a, 1b and 1c, for Majorana neutrinos we have plotted the parameter space corresponding to any of the two mixing angles by constraining the third angle by its values given in equation (15) while giving full allowed variation to other parameters. Also included in the figures are blank rectangular regions indicating the experimentally allowed 3σ region of the plotted angles. Interestingly, a general look at these figures reveals that the case of inverted hierarchy seems to be ruled out. From figure 1a showing the plot of angles θ_{12} versus θ_{23} , one can immediately conclude that the plotted parameter space includes the experimentally allowed range of $\theta_{23} = 35.7^\circ - 55.6^\circ$, however it excludes the experimentally allowed range of $\theta_{12} = 29.3^\circ - 39.25^\circ$. This clearly indicates that at 3σ C.L. inverted hierarchy is not viable. The conclusions arrived above can be further checked from figures 1b and 1c wherein we have plotted θ_{12} versus θ_{13} and θ_{23} versus θ_{13} respectively by constraining angles θ_{23} and θ_{12} . Without getting into the details, one can show that the degenerate scenarios mentioned above are also ruled out for these mass matrices. Further, from the above analysis of texture 4 zero mass matrices,




Figure 1: Plots showing the parameter space corresponding to any of the two mixing angles by constraining the third angle by its experimental limits given in equation (15) and giving full allowed variation to other parameters for Majorana neutrinos. The blank rectangular region indicates the experimentally allowed 3σ region of the plotted angles.

one can easily deduce similar conclusions for texture 6 zero as well as for the two cases of the texture 5 zero mass matrices.

Coming to the normal hierarchy cases, detailed dependence of mixing angles on the lightest neutrino mass as well as the parameter space available to the phases of mass matrices have been investigated for both Majorana and Dirac neutrinos [17, 18]. Without presenting the details, several phenomenological quantities such as the viable ranges of neutrino masses, mixing angle s_{13} , Jarlskog's rephasing invariant parameter J_l , the Dirac-like CP violating phase δ_l and the effective neutrino mass $\langle m_{ee} \rangle$, related to neutrinoless double beta decay $(\beta\beta)_{0\nu}$, have also been calculated for different cases. In the case of texture 4 zero mass matrices, for Majorana neutrinos, we obtain $m_{\nu_1} = 2.47 \times 10^{-4} - 0.006$, $\theta_{13} = 1.14^\circ - 11.50^\circ$ and $J_l = -0.0459 - .0463$, whereas for Dirac neutrinos, these come out to be $m_{\nu_1} = 5.73 \times 10^{-5} - 0.012$, $\theta_{13} = 0.084^\circ - 11.50^\circ$ and $J_l = -0.0462 - .0448$. Our analysis reveals that a measurement of m_{ν_1} could have important implications for the nature of neutrinos. Also, the lower limit on θ_{13} for the Dirac case is considerably lower than for the Majorana case, therefore a measurement of θ_{13} would have important implications for this case.

For the sake of completion, we have also carried out calculations wherein the charged lepton mass matrix is in the flavor basis for all the texture specific cases considered here. As expected, the inverted hierarchy and degenerate scenarios are ruled out for all the Dirac and Majorana cases. For the normal hierarchy, the texture 6 zero and the texture 5 zero $D_\nu = 0$ case are again ruled out. For the texture 5 zero $D_l = 0$ case and the texture 4 zero mass matrices, the ranges of various phenomenological quantities calculated here are much narrower compared to the corresponding texture specific cases.

To summarize, implications of Fritzsch-like Hermitian texture specific mass matrices have been investigated for both quarks and neutrinos. In the case of quarks, our analysis reveals that texture 4 zero mass matrices are quite compatible with well known parameter $\sin 2\beta$ and with other precise CKM matrix elements. In the case of leptonic mass matrices,

for Majorana and Dirac neutrinos, we find that for texture 4, 5, 6 zero mass matrices the inverted hierarchy and degenerate scenarios of neutrino masses are ruled out by the present data.

Acknowledgements

GA would like to thank DST, Government of India for financial support and the Chairman, Department of Physics for providing facilities to work. RV would like to thank the Director, RIEIT, for providing facilities to work.

References

- [1] H. Fritzsch, Z. Z. Xing, Prog. Part. Nucl. Phys. 45, (2000) 1, and references therein.
- [2] H. Fritzsch, Z. Z. Xing, Prog. Part. Nucl. Phys. J. C36, (2004) 227, and references therein.
- [3] N. Cabibbo, Phys. Rev. Lett. 10, (1963) 531; M. Kobayashi, T. Maskawa, Prog. Theor. Phys. 49, (1973) 652.
- [4] C. Amsler *et al.*, Phys. Lett. B667, (2008) 1, updated results available at <http://pdg.lbl.gov/>.
- [5] G. Ahuja, M. Gupta, S. Kumar, M. Randhawa, Phys. Lett. B647, (2007) 394.
- [6] G. L. Fogli, E. Lisi, A. Marrone, A. Melchiorri, A. Palazzo, A. M. Rotunno, P. Serra, J. Silk, A. Slosar, Phys. Rev. D78, (2008) 033010; G. L. Fogli, E. Lisi, A. Marrone, A. Palazzo, A. M. Rotunno, arXiv:0809.2936.
- [7] R. G. Roberts, A. Romanino, G. G. Ross, L. Velasco-Sevilla, Nucl. Phys. B615, (2001) 358.
- [8] H. Fritzsch, Z. Z. Xing, Phys. Lett. B555, (2003) 63.
- [9] H. D. Kim, S. Raby, L. Schradin, Phys. Rev. D69, (2004) 092002.
- [10] Z. Z. Xing, H. Zhang, J. Phys. G30, (2004) 129.
- [11] K. Matsuda, H. Nishiura, Phys. Rev. D74, (2006) 033014.
- [12] Z. Z. Xing, H. Zhang, S. Zhou, Phys. Rev. D77, (2008) 113016; H. Leutwyler, Phys. Lett. B378, (1996) 313.
- [13] J. Charles *et al.*, CKMfitter Group, Eur. Phys. J. C41, (2005) 1, updated results available at <http://ckmfitter.in2p3.fr/>.
- [14] M. Bona *et al.*, UTfit Collaboration, JHEP 0610, (2006) 081, updated results available at <http://www.utfit.org/>.

- [15] E. Barberio *et al.*, Heavy Flavor Averaging Group (HFAG), hep-ex/0603003, updated results available at <http://www.slac.stanford.edu/xorg/hfag/>.
- [16] M. Randhawa, G. Ahuja, M. Gupta, Phys. Rev. D65, (2002) 093016.
- [17] M. Randhawa, G. Ahuja, M. Gupta, Phys. Lett. B643, (2006) 175.
- [18] G. Ahuja, S. Kumar, M. Randhawa, M. Gupta, S. Dev, Phys. Rev. D76, (2007) 013006.